


# Pooled sputum testing for molecular TB diagnosis – Lessons from the Start4All multi-country experience

LabCoP Extended ECHO Session  
16<sup>th</sup> April 2026



# Start4All





**Start4All**

# Objectives of the webinar

Dr Tom Wingfield

Start4All Chief-Investigator

[tom.wingfield@lstmed.ac.uk](mailto:tom.wingfield@lstmed.ac.uk)

<https://lstmed.ac.uk/people/tom-wingfield/>

[www.linkedin.com/in/tom-wingfield-46716a339/](http://www.linkedin.com/in/tom-wingfield-46716a339/)

# Objectives



**Start4All**

- Describe the **concept of pooled sputum testing** for molecular TB diagnosis, including when it may be appropriate for routine laboratory practice.
- Summarize **key findings from the Start4All 7-country study** on diagnostic performance, cartridge efficiency, and operational feasibility.
- Identify **practical laboratory considerations** for implementation, including workflow adaptation, use of existing equipment and personnel, and quality-assured testing in routine settings.
- Discuss how pooled testing may **support laboratory network efficiency**, decentralization of TB diagnosis, and future implementation priorities.



**Start4All**

# Overview of Start4All

# The Start4All Partnership



# Start4All: Start taking action for TB diagnosis




**Start4All**

A global project evaluating screening and diagnostic algorithms for TB



**Professor Luis Cuevas**





### Phase 1

Evaluate the performance, costs, feasibility, and acceptability of *selected* TB screening & diagnostic tests and test algorithms at:

- the lowest levels of health care (primary healthcare and district hospitals)
- amongst key and underserved populations in community settings.



2024 -  
2025

# Participants



**Start4All**

**14 741** participants were recruited during Phase 1

Facility-based case finding (**9349**)

*Bangladesh, Brazil, Cameroon, Kenya, Malawi, Nigeria, Viet Nam*



Primary Healthcare Centres (PHC) 4409



District Hospital outpatient clinics (DH) 4134

Community-based case finding (**5392**)



Nomads  
*Nigeria*



IDP, informal settlements, rural and urban vulnerable  
*Bangladesh, Cameroon, Nigeria, & Viet Nam*

Children (**806**)



Predominantly facility-based  
*Cameroon and Viet Nam*


# Tests and approaches evaluated




**Start4All**



Xpert Ultra *pooled*  
testing on sputum  
(1:4 but allow 1:2 & 1:3)



CRP



FujiLAM assay



CAD CXR



Xpert Ultra  
Index test



Culture (MGIT / LJ)

Reference standard

# Start4All Phase 1 Take Aways



**Start4All**

- Strong performance of CAD-CXR & Pooled Xpert Ultra algorithms
- Pooled testing showed significant cartridge and cost savings but with some decrease in detection (~3% lower sensitivity)
- Pooled TB testing could enhance equitable access (if savings redistributed)



**World Health  
Organization**

## Recommendation


**When resource constraints do not allow for the testing of individual samples:** In adults and adolescents with signs and symptoms for pulmonary TB or who screen positive for pulmonary TB, LC-aNAATs on (up to 4) pooled sputa may be used as the initial diagnostic strategy for diagnosing TB rather than LC-aNAATs on individual samples (*conditional recommendation, high certainty of evidence*).

- Start4All team implementing & evaluating best algorithms in Phase 2

# Start4All Pooled Testing Toolkit



**Start4All**



A practical toolkit to support safe, efficient pooled sputum testing.

**Who it's for:** National TB programmes, laboratories, clinicians, and implementation teams.

**What's included:**

- Standard operating procedures (SOPs)
- FAQs and site readiness checklist
- Two case studies
- Laboratory poster
- Training presentation
- Multilingual videos



Available **free** online now at [www.lstmed.ac.uk/start4all/toolkits](http://www.lstmed.ac.uk/start4all/toolkits)